

MSC: 42A32, 33B30, 41A10, 11M06, 33B15

DOI: 10.21538/0134-4889-2022-28-4-177-190

**UNIFORM WITH RESPECT TO THE PARAMETER  $a \in (0, 1)$  tWO-SIDED ESTIMATES OF THE SUMS OF SINE AND COSINE SERIES WITH COEFFICIENTS  $1/k^a$  BY THE FIRST TERMS OF THEIR ASYMPTOTICS**

**A. Yu. Popov, T. V. Rodionov**

Uniform with respect to the parameter  $a \in (0, 1)$  estimates of the functions  $f_a(x) = \sum_{k=1}^{\infty} k^{-a} \cos kx$  and  $g_a(x) = \sum_{k=1}^{\infty} k^{-a} \sin kx$  by the first terms of their asymptotic expansions  $F_a(x) = \sin(\pi a/2)\Gamma(1-a)x^{a-1}$  and  $G_a(x) = \cos(\pi a/2)\Gamma(1-a)x^{a-1}$  are obtained. Namely, it is proved that the inequalities

$$G_a(x) - \frac{x}{2} < g_a(x) < G_a(x) - \frac{x}{12}$$

and

$$F_a(x) + \zeta(a) + \frac{\zeta(3)}{4\pi^3} x^2 \sin(\pi a/2) < f_a(x) < F_a(x) + \zeta(a) + \frac{1}{18} x^2 \sin(\pi a/2).$$

are valid for all  $a \in (0, 1)$  and  $x \in (0, \pi]$ .

It is shown that the estimates are unimprovable in the following sense. In the lower estimate for the sine series, the subtrahend  $x/2$  cannot be replaced by  $kx$  with any  $k < 1/2$ : the estimate ceases to be fulfilled for sufficiently small  $x$  and the values of  $a$  close to 1. In the upper estimate, the subtrahend  $x/12$  cannot be replaced by  $kx$  with any  $k > 1/12$ : the estimate ceases to be fulfilled for the values of  $a$  and  $x$  close to 0. In the lower estimate for the cosine series, the multiplier  $\zeta(3)/(4\pi^3)$  of  $x^2 \sin(\pi a/2)$  cannot be replaced by any larger number: the estimate ceases to be fulfilled for  $x$  and  $a$  close to 0. In the upper estimate for the cosine series, the multiplier  $1/18$  of  $x^2 \sin(\pi a/2)$  can probably be replaced by a smaller number but not by  $1/24$ : for every  $a \in [0.98, 1)$ , such an estimate would not hold at the point  $x = \pi$  as well as on a certain closed interval  $x_0(a) \leq x \leq \pi$ , where  $x_0(a) \rightarrow 0$  as  $a \rightarrow 1$ -. The obtained results allow us to refine the estimates of the functions  $f_a$  and  $g_a$  established recently by other authors.

Keywords: special trigonometric series, polylogarithm, periodic zeta function.

**REFERENCES**

1. Bieberbach L. *Analytische Fortsetzung*. Berlin: Springer-Verlag, 1955, 168 p. doi: 10.1007/978-3-662-01270-3. Translated to Russian under the title *Analiticheskoe prodolzhenie*. Moscow: Nauka Publ., 1967, 240 p.
2. Zygmund A. *Trigonometric series*, vol. I, II. Cambridge: Cambridge Univ. Press, 1959; vol. I, 383 p.; vol. II, 354 p. Translated under the title *Trigonometricheskie ryady*, M.: Mir Publ., 1965, vol. I, 616 p; vol. II, 538 p.
3. Titchmarsh E.C. *The theory of the Riemann zeta-function*. Oxford: Oxford Univ. Press, 1987, 422 p. ISBN: 0198533691. Translated to Russian under the title *Teoriya dzeta-funktsii Rimana*, Moscow: Izd. Inostr. Lit., 1953, 407 p.
4. Erdélyi A. (ed.) *Higher transcendental functions*. Vol. 1. NY: McGraw Hill, 1953, 302 p.
5. Leau L. Recherches des singularités d'une fonction définie par un développement de Taylor. *Journ. de Math. (5)*, 1899, vol. 5, pp. 365–425.
6. Lifyand E., Podkorytov A. Lebesgue constants of Riesz type means of negative order. *J. Math. Anal. Appl.*, 2022, vol. 505, no. 2, art. no. 125618. doi: 10.1016/j.jmaa.2021.125618.
7. Lindelöf E.L. *Le calcul des résidus et ses applications à la théorie des fonctions*. Paris: Gauthier–Villar, 1905, 158 p.
8. Olver F.W.J. et al. (eds.) *NIST handbook of mathematical functions*. NY: Cambridge Univ. Press, 2010, 968 p. The online version: The NIST Digital Library of Mathematical Functions (DLMF): <https://dlmf.nist.gov/>.

9. Truesdell C. On a function which occurs in the theory of the structure of polymers. *Ann. Math. (2)*, 1945, vol. 46, no. 1, pp. 144–157. doi: 10.2307/1969153.

Received May 19, 2022

Revised July 29, 2022

Accepted August 4, 2022

**Funding Agency:** The research of the first author (the results of Sections 2–3) was carried out at Moscow State University and supported by the Russian Science Foundation (project no. 22-21-00545). The research of the second author (the results of Section 6) was carried out at Moscow State University and supported by the Russian Foundation for Basic Research (project no. 20-01-00584).

*Anton Yur'evich Popov*, Dr. Phys.-Math. Sci., Lomonosov Moscow State University and Moscow Centre of Fundamental and Applied Mathematics, Moscow, 119991 Russia, e-mail: station@list.ru.

*Timofey Victorovich Rodionov*, Cand. Sci. (Phys.-Math.), Lomonosov Moscow State University and Moscow Centre of Fundamental and Applied Mathematics, Moscow, 119991 Russia, e-mail: rodionovtv@mail.ru.

Cite this article as: A. Yu. Popov, T. V. Rodionov. Uniform with respect to the parameter  $a \in (0, 1)$  two-sided estimates of the sums of sine and cosine series with coefficients  $1/k^a$  by the first terms of their asymptotics. *Trudy Instituta Matematiki i Mekhaniki UrO RAN*, 2022, vol. 28, no. 4, pp. 177–190.